INFLIXIMAB PRESCRIBER ORDER FORM

Fax completed form, insurance information, and clinical documentation to:

Patient Name:			**Date of Birth:**	
Address:				
Phone:	**Height:**	☐ inches ☐ cm	**Weight:**	☐ lbs ☐ kg

Clinical Information

Primary Diagnosis Description:		**ICD-10 Code:**	
Is this the first dose?	☐ Yes – date of first dose: ☐ No – date of next dose due:	**Hepatitis B Status:**	Titer Date: ☐ Positive ☐ Negative
TB Status:	☐ PPD (negative) – date: ☐ Last chest x-ray – date: ☐ Past positive TB infection, course taken:	☐ Active TB ☐ Unknown	

Infliximab Prescription

☐ **Infliximab (Remicade®) *or* ☐ Infliximab-dyyb (Inflectra®) *or* ☐ Infliximab-abda (Renflexis®) refill as directed x 1 year**

Initial Dose: ☐ Infuse _____ mg/kg IV on Weeks 0, 2, and 6.

☐ Other: ______________________________

Maintenance Dose: ☐ Infuse _____ mg/kg IV every 8 weeks.

☐ Other: ______________________________

Dose will be rounded to closest 100 mg vial.

Infusion will be given at a flat rate over 2 hours unless patient has a history of infusion-related reaction(s), and then will infuse with a titrated rate.

Ancillary Orders

Anaphylaxis Kit

If this is a 1st dose, would you like Option Care Health to provide an anaphylaxis kit with the 1st dose?

☐ Yes – please complete Anaphylaxis Physician Order (FR-PC-036) provided ☐ No

Medication Orders

- ☐ Acetaminophen 650 mg PO 30 min before infusion, may repeat every 3 to 4 hours as needed for fever or mild discomfort. Patient may use own supply or patient may decline.
- ☐ Diphenhydramine 25 mg PO 30 min before infusion, may repeat every 4 to 6 hours as needed for mild to moderate allergic reactions. Patient may use own supply or patient may decline.
- ☐ Loratadine 10 mg PO 30 min before infusion. Patient may use own supply or patient may decline.
- ☐ Methylprednisolone 40 mg IV push 20 minutes prior to infusion.
- ☐ Other: ______________________________

IV Flush Orders

- ☐ Peripheral: NS 2 to 3 mL pre-/post-use.
- ☐ Implanted Port: NS 5 to 10 mL pre-/post-use and 10 to 20 mL pre-/post-lab draw. Heparin (100 unit/mL) 3 to 5 mL post-use. For maintenance, heparin (100 unit/mL) 3 to 5 mL every 24 hr if accessed or weekly to monthly if not accessed.

Lab Orders

- ☐ No labs ordered at this time.
- ☐ Other: ______________________________

Skilled nurse to administer doses intravenously in the home or alternate care setting. Refill above ancillary orders as directed x 1 year.

I certify that the use of the indicated treatment is medically necessary and I will be supervising the patient's treatment.

Prescriber Signature: ______________________________ **Date:** ______________

Prescriber Information

Prescriber Name:		**Phone:**	**Fax:**
Address:		**NPI:**	
City, State:	**Zip:**	**Office Contact:**	

CONFIDENTIAL HEALTH INFORMATION: Healthcare information is personal information related to a person's healthcare. It is being faxed to you after appropriate authorization or under circumstances that do not require authorization. You are obligated to maintain it in a safe, secure, and confidential manner. Re-disclosure of this information is prohibited unless permitted by law or appropriate customer/patient authorization is obtained. Unauthorized re-disclosure or failure to maintain confidentiality could subject you to penalties described in federal and state laws. **IMPORTANT WARNING**: This message is intended for the use of the person or entity to whom it is addressed and may contain information that is privileged and confidential, the disclosure of which is governed by applicable law. If the reader of this message is not the intended recipient, or the employee or agent responsible for delivering it to the intended recipient, you are hereby notified that any dissemination, distribution or copying of this information is STRICTLY PROHIBITED. If you have received this message in error, please notify us immediately. Brand names are the property of their respective owners.

 @2020 UBACARE. All Rights Reserved.